

Research article

# Transpiration, CO<sub>2</sub> assimilation, WUE, and stomatal aperture in leaves of *Viscum album* (L.): Effect of abscisic acid (ABA) in the xylem sap of its host (*Populus x euamericana*)

Peter Escher<sup>a,\*</sup>, Andreas D. Peuke<sup>a</sup>, Peter Bannister<sup>b</sup>, Siegfried Fink<sup>c</sup>,  
Wolfram Hartung<sup>d</sup>, Fan Jiang<sup>e</sup>, Heinz Rennenberg<sup>a</sup>

<sup>a</sup> Institute of Forest Botany and Tree Physiology, Chair of Tree Physiology, Albert-Ludwigs-University Freiburg,  
Georges-Köhler-Allee 053/054, D-79110 Freiburg i.B., Germany

<sup>b</sup> Department of Botany, University of Otago, P.O. Box 56, Dunedin, New Zealand

<sup>c</sup> Institute of Forest Botany and Tree Physiology, Chair of Forest Botany, Albert-Ludwigs-University Freiburg,  
Bertoldstrasse 17, 79085 Freiburg i.B., Germany

<sup>d</sup> Julius-von-Sachs-Institut für Biowissenschaften, Lehrstuhl für Botanik I, Julius-Maximilians Universität Würzburg,  
Julius-von-Sachs-Platz 2, D-97082 Würzburg, Germany

<sup>e</sup> Institute of Life Sciences, Beijing Normal University, Beijing, China

Received 11 December 2006; accepted 25 July 2007

Available online 31 July 2007

## Abstract

Leaves of the mistletoe *Viscum album* (L.) show a high rate of transpiration, even when the host is under severe drought stress. The hypothesis that a strong control of ABA influx from the xylem sap of the host into the mistletoe prevents stomatal closure in mistletoe leaves was tested under the following conditions: sections of poplar twigs carrying a mistletoe were perfused with artificial xylem sap that contained different ABA concentrations and both transpiration and ABA levels were analysed in mistletoe leaves. Despite variation by a factor of 10<sup>4</sup>, the ABA content of the host xylem did not affect ABA levels, leaf transpiration, CO<sub>2</sub> assimilation, WUE, or the degree of stomatal aperture in mistletoe leaves. These observations support the hypothesis of a strong control of ABA influx from the host of the xylem into the mistletoe, although degradation of ABA before it enters the mistletoe leaves cannot be excluded. This mechanism may ensure a water and nutritional status favourable for the mistletoe, even if the water status of the host is impaired.

Despite the lack of short-term sensitivity of ABA levels in mistletoe leaves to even strong changes of ABA levels in the xylem sap of the host, ABA levels in mistletoe leaves were relatively high compared to ABA levels in the leaves of several tree species including poplar. Since significant transpiration of the mistletoe leaves was observed despite high ABA levels, a diminished sensitivity of the stomata of mistletoe leaves to ABA has to be concluded. The stomatal density of adaxial *Viscum* leaves of 89 ± 23 stomata per mm is lower than those reported in a study performed at the end of the 19th century.

© 2007 Published by Elsevier Masson SAS.

**Keywords:** ABA; Abscisic acid; Assimilation; Stomatal aperture; Transpiration; Xylem sap

## 1. Introduction

One of the most striking features of the mistletoe *Viscum album* (L.) is its high rate of transpiration, even when the

host is under severe drought stress [23,54]. As a consequence, twigs of the host that are distal to the insertion of a mistletoe often dry out completely and die [53]. For several reasons, this observation is surprising. Excessive water loss of leaves under drought stress is usually prevented to some extent by regulating the degree of stomatal aperture [40,52] and abscisic acid (ABA) produced in the roots and allocated to the leaves with the transpiration stream is a central component of the

\* Corresponding author. Tel.: +49 (0) 761 203 8300; fax: +49 (0) 761 203 8302.

E-mail address: [peter.escher@ctp.uni-freiburg.de](mailto:peter.escher@ctp.uni-freiburg.de) (P. Escher).

signalling cascade that leads to stomatal closure [16,25]. *V. album* does not rely only on the water supplied by the xylem sap of its host, but also uses the xylem sap of its host as a source of carbon, nitrogen, and sulfur for growth and development [18,21,19,20]. Therefore, it would be expected that changes in the allocation of ABA in the xylem sap of the host would also regulate the stomatal aperture of mistletoe leaves. However this is obviously not the case and may be explained by (i) a very effective control of ABA influx from the host xylem into the mistletoe, or (ii) an ABA insensitivity of the stomata of mistletoe leaves.

Early studies suggested that *V. album* lacked ABA [22,27] and was later disproved by Ihl et al. [29], who found low ABA levels in July and high levels in wintertime. Since low ABA levels in mistletoe leaves coincide with low rates of transpiration and low levels of ABA with high rates of transpiration [29], ABA insensitivity of mistletoe leaves does not seem to be responsible for the lack of stomatal closure under conditions of drought stress. The aim of the present experiments was to test the hypothesis that a strong control of ABA influx from the xylem sap of the host into the mistletoe prevents stomatal closure under drought stress in mistletoe leaves. Consequently, sections of poplar twigs bearing a mistletoe were perfused with artificial xylem sap that contained different ABA concentrations and both gas exchange and ABA levels were analysed in mistletoe leaves.

## 2. Methods

### 2.1. Plant material

The present experiments were performed with twig sections of *Populus x euamericana* carrying a mistletoe (*V. album* L.). Plant material was collected in the upper Rhine valley at a site near Rheinstetten (about 10 km south-west of Karlsruhe, Germany; co-ordinates 48°98'N, 08°33'E) in a 60–70 years old riparian lowland forest of *P. x euamericana* on the banks of a small water channel. Long-term annual air temperature in this region is 10.3 °C and annual precipitation is 770 mm [38]. On 22nd August 2005, a total of 20 twigs were collected with a lifting platform between 10 a.m. and 2 p.m. from three trees. The cut ends of the twigs were immediately transferred into a bucket filled with water to avoid embolism of the vessels.

### 2.2. Perfusion of poplar twigs

Poplar twigs carrying a mistletoe were perfused with artificial xylem sap [21]. For this purpose, twigs were cut off at both ends, resulting in a 10–15 cm section with a mistletoe attached. Approximately 2–3 cm bark was immediately peeled off at the proximal end and the wood was attached to the perfusion system as previously described [21]. Subsequently, the perfusion solution was pumped continuously through the twig for 20 h. In order to achieve close to natural osmotic conditions and a close to natural nutrient status, the perfusion solution contained 2 mM L-Gln, 2 mM K<sup>+</sup>, 0.5 mM Mg<sup>2+</sup>, 0.75 mM Ca<sup>2+</sup>,

0.75 mM H<sub>2</sub>PO<sub>4</sub><sup>-</sup>, 0.5 mM SO<sub>4</sub><sup>2-</sup>, 1.75 mM Cl<sup>-</sup>. The pH was adjusted to 6.5 according to Refs. [7,8] to mimic the natural pH of the host xylem sap, to prevent dissociation of ABA and to keep ABA permeable across membranes. Solutions containing 0.01 μM (±)-*cis/trans*-abscisic acid (MoBi Tec, Göttingen, Germany) were applied for the initial perfusion for 3 h. Subsequently, the initial perfusion solution was replaced by solutions, which contained ABA at four different concentrations (0.01, 0.1, 1, 100 μM). Experiments were performed in a controlled climate chamber (Heräeus Vötsch HPS 1500, Hanau, Germany) at 25 °C, 50 ± 2% relative humidity, and ca. 430 μE m<sup>-2</sup> s<sup>-1</sup> PPFD (provided by Philips TLD 58 W/550 and OSRAM L58 W/77 Fluora lamps).

### 2.3. Transpiration and CO<sub>2</sub> assimilation of mistletoe leaves

Transpiration and CO<sub>2</sub> assimilation of individual pairs of mistletoe leaves were measured at 3, 6, 13, and 20 h after beginning of the experiment with a GFS-3000 portable gas exchange fluorescence system connected to a 3010-S standard measuring head with a 3040-L LED light source (Heinz Walz GmbH, Effeltrich, Germany). In order to assess the specific transpiration (*E*) and CO<sub>2</sub> assimilation (*A*) of the mistletoe, the individual leaf areas were determined with an area meter (Δ*T* area meter, Delta-T Devices, Cambridge, UK). Water use efficiency (WUE) was calculated by the following equation:

$$\text{WUE} = A \text{ [mmol CO}_2 \text{ m}^{-2} \text{ s}^{-1}] / E \text{ [mmol H}_2\text{O m}^{-2} \text{ s}^{-1}]$$

### 2.4. Stomatal density and degree of stomatal opening

The upper adaxial surface of one leaf of each mistletoe was coated with UHU Hart (UHU GmbH & Co. KG, Bühl, Germany) immediately after the experiment. The cured glue conserved imprints of the leaf surfaces, enabled the analysis of the degree of stomatal opening at the end of experiment, and restricted possible shrinking effects [28]. Subsequently, hardened glue was observed with a microscope using the differential interference contrast techniques (DIC) (Zeiss Axiophot, equipped with a CCD-camera Axiocam MRc 5). After focusing, five pictures of each leaf were taken at 20 times magnification with 1292 × 968 pixel and saved as 8 bit greyscale tif files (Fig. 2). Subsequent image analyses were performed with ImageJ [47].

For the determination of stomatal density, one picture of each leaf was used. Stomata were identified and marked manually to obtain the number of stomata per frame and the spatial distribution (2D) data. To normalise the results and to make data comparable, euclidian distances were calculated between stomata to access mean euclidian distance as a leaf parameter. To describe stomatal distribution statistically [24], the degree to which the observed pattern departs from a random pattern [*R*] was calculated according to Ref. [13] with the limits *R* = 0, i.e. aggregated pattern individuals are

at same locus, distance to the nearest neighbour is 0;  $R = 2.1491$ , i.e. perfectly uniform pattern with maximal hexagonal spacing; and  $R = 1$ , i.e. pattern with random distribution.

For determination of stomatal aperture, the perimeters of plainly visible stomata were marked manually in five pictures per leaf. Subsequently, area, bounding rectangle perimeter, and minor and major axis of a fitting ellipse of the stomata were measured simultaneously with ImageJ [47]. Due to the large variation of the lengths of stomata, the area of aperture is not directly comparable. To avoid these difficulties, “relative stomatal aperture (RSA)” according to Ref. [32] as a size independent variable was calculated. RSA [%] is specified as  $(400 \times A)/(\Pi \times L^2)$ , where  $A$  is the aperture area and  $L$  is the length of pore  $\approx$  major axis of a fitting ellipse.

### 2.5. Determination of ABA concentrations in leaves

Three leaves were cut from each mistletoe before and directly after the experiment, immediately frozen with liquid nitrogen, and stored at  $-80^\circ$  until further analysis. For determination of ABA content, leaves were lyophilised, ground with mortar and pestle, and aliquots of the three leaves of each mistletoe were cumulated. Freeze-dried tissue samples were homogenised and extracted in 80% aqueous methanol. Extracts were passed through a Sep Pak C<sub>18</sub>-cartridge. Methanol was removed under reduced pressure and the aqueous residue was partitioned three times against ethyl acetate at pH 3.0. The ethyl acetate of the combined organic fractions was removed under reduced pressure. The newly obtained residue was taken up in TBS-buffer (tris buffered saline; 150 mmol/L NaCl, 1 mmol/L MgCl<sub>2</sub> and 50 mmol/L tris at pH 7.8) and subjected to an immunological ABA assay (ELISA) as described earlier [37,43]. The accuracy of the ELISA has been verified in earlier investigations [43]. Recoveries of ABA during the purification procedures were checked routinely using radioactive ABA and found to be higher than 95%. The immunochemicals were generously supplied by Prof. Weiler, Ruhr Universität Bochum (Germany).

### 2.6. Data analysis

Transpiration rates of one selected pair of leaves of each of the 20 mistletoes were measured repeatedly at 3, 6, 13, 20 h. One mistletoe, which did not show any significant transpiration during the experiment, was discarded from further data analysis as we did also for severe (negative) outliers in assimilation and WUE. Analysis of variance was done by GLM-ANOVA with Tukey–Kramer multiple-comparison post hoc test. All statistical analyses were carried out with NCSS 2004 [26]. In boxplots, the top and the bottom of the box represent the 25th and 75th percentile. The length of the box is equal to the interquartile range (*IQR*). Lines drawn in the box represent the median. T-shaped lines are the adjacent values  $\leq 75\%$  ( $\geq 25\%$ ) percentile + 1.5 *IQR* ( $-1.5$  *IQR*).

## 3. Results

### 3.1. Transpiration, assimilation, and WUE of mistletoes at increasing ABA concentrations in the host xylem

Specific transpiration of mistletoe leaves ( $E$ ) was highest in controls at 3 h exposure ( $2.5 \pm 1.2$  mmol H<sub>2</sub>O m<sup>-2</sup> h<sup>-1</sup>) (Fig. 1) and subsequently declined significantly to  $0.8 \pm 0.4$  mmol H<sub>2</sub>O m<sup>-2</sup> h<sup>-1</sup> at 13 h exposure. But there was no significant effect of the ABA concentration in the xylem sap of the host ranging from 0.01 to 100  $\mu$ M, on transpiration of the mistletoe leaves during the entire experiment (Fig. 1). Assimilation rates ( $A$ ) of the mistletoe leaves ranged from 2.3 to 10.2 mmol CO<sub>2</sub> m<sup>-2</sup> s<sup>-1</sup> with an overall mean of  $4.5 \pm 3.5$  mmol CO<sub>2</sub> m<sup>-2</sup> s<sup>-1</sup> (Fig. 2). There were no significant differences due to the time of exposure or the different ABA concentrations in the xylem sap of the host. The WUE of C assimilation was calculated from the individual ratio between  $A$  and  $E$  for each mistletoe; it increased in controls from 3 to 6 h of exposure, then remained constant, and was not affected by the ABA concentration in host xylem sap (Fig. 3).

### 3.2. ABA concentrations in leaf material

ABA concentrations in mistletoe leaves ranged between 6.9 and 31.8  $\mu$ mol g dw<sup>-1</sup> (Fig. 6). Despite the wide range of ABA concentrations applied, the ABA concentration in the mistletoe leaves was not significantly affected by the ABA concentration in the xylem sap of the host.




Fig. 1. Transpiration rate  $E$  (mmol m<sup>-2</sup> s<sup>-1</sup>) of the leaves of mistletoes exposed to different ABA levels via the xylem sap of the host.  $E$  was measured at 3, 6, 13 and 20 h after starting the perfusion experiment at 25 °C and 50% relative humidity. In controls, 0.01  $\mu$ M ABA was fed into host xylem during the entire experiment. In the other series, the perfusion solution containing 0.01  $\mu$ M ABA was replaced after 3 h by solutions containing 0.1, 1 or 100  $\mu$ M ABA. Significant differences in  $E$  between treatments (ABA levels) were not observed at any individual time. \* Indicates significant differences in controls after different times of perfusion.




Fig. 2. Assimilation rate  $A$  ( $\mu\text{mol m}^{-2} \text{s}^{-1}$ ) of the leaves of mistletoes exposed to different ABA levels via the xylem sap of the host.  $A$  was measured at 3, 6, 13 and 20 h after starting the perfusion experiment at 25 °C and 50% relative humidity. In controls, 0.01  $\mu\text{M}$  ABA was fed into host xylem during the entire experiment. In the other series, the perfusion solution containing 0.01  $\mu\text{M}$  ABA was replaced after 3 h by solutions containing 0.1, 1 or 100  $\mu\text{M}$  ABA. Significant differences in  $A$  between treatments (ABA levels) at any individual time or at the same treatment after different times of perfusion were not observed.

### 3.3. Stomatal density and stomatal aperture

Mean stomatal density of *Viscum* leaves (Fig. 4) was  $89 \pm 23$  stomata per  $\text{mm}^{-2}$  with a range between 36 and 121 stomata




Fig. 3. Water use efficiency (WUE) of the leaves of mistletoes exposed to different ABA levels via the xylem sap of the host. WUE was calculated at 3, 6, 13 and 20 h after starting the perfusion experiment at 25 °C and 50% relative humidity. In controls, 0.01  $\mu\text{M}$  ABA was fed into host xylem during the entire experiment. In the other series, the perfusion solution containing 0.01  $\mu\text{M}$  ABA was replaced after 3 h by solutions containing 0.1, 1 or 100  $\mu\text{M}$  ABA. Significant differences in WUE between treatments (ABA levels) were not observed at any individual time. \* Indicates significant differences in controls after different times of perfusion.




Fig. 4. Gray-scale image of stomata on the leaf surface of *Viscum album*. The images were obtained by light microscopy of glue imprints. The white bar represents a distance of 400  $\mu\text{m}$ . For measuring stomatal aperture, pictures with a 20-fold higher magnification to this picture were used (for example see the not scaled inset in the left up corner).

per  $\text{mm}^{-2}$  (Table 1). Mean euclidian distance (ED) varied between 352 and 446  $\mu\text{m}$ .  $R$  calculated with minimum ED and the number of stomata per unit area showed a mean value of  $1.53 \pm 0.13$ , indicating that stomatal pattern is about half between random distribution and perfectly uniform hexagonal spacing. Despite the broad range in the observed leaf parameters, outliers were not found (Mahalanobis distance) and the leaf material of the examined mistletoe specimen was relatively homogenous. Relative stomatal aperture (RSA) ranged between 12.4% and 14.2% and was not significantly affected by the concentration of ABA supplied in the xylem of the host (Fig. 5).

## 4. Discussion

The present perfusion experiments show that differences in the ABA content of the host xylem do not cause short-term effects in mistletoe leaves on ABA levels, leaf transpiration,  $\text{CO}_2$  assimilation, WUE, or the degree of stomatal aperture. For two reasons this result is surprising. On one hand, the ABA concentrations were chosen to simulate water stress [12] and should induce a fast [11,35] stomatal response in plant leaves [3,7]. On the other hand, the ABA concentrations in the xylem sap of the host were varied 10<sup>4</sup>-fold and were several orders of magnitude higher than ABA concentrations reported for the xylem sap (0.2–2.2  $\mu\text{M}$ ) of *Populus* species

Table 1

Parameters of the spatial distribution of stomata on the adaxial side of mistletoe leaves (median euclidian distance, ED; mean euclidian distance; minimum euclidian distance; maximum euclidian distance; stomatal density,  $n$  ( $\text{mm}^{-2}$ ); and  $R$ )

| Median ED ( $\mu\text{m}$ ) | Mean ED ( $\mu\text{m}$ ) | Min. ED ( $\mu\text{m}$ ) | Max. ED ( $\mu\text{m}$ ) | $n$ ( $\text{mm}^{-2}$ ) | $R$             |
|-----------------------------|---------------------------|---------------------------|---------------------------|--------------------------|-----------------|
| 411 $\pm$ 19                | 424 $\pm$ 19              | 42 $\pm$ 16               | 996 $\pm$ 59              | 89 $\pm$ 23              | 1.53 $\pm$ 0.12 |






Fig. 5. “Relative stomatal aperture (RSA)” [%] of mistletoe leaves after 20 h perfusion of the host xylem with different ABA concentrations. Significant differences in RSA between treatments (ABA levels) were not observed.

[11,10]. Apparently, short-term fluctuations in ABA production by the host roots, its allocation to and action in the host leaves, and drought stress, do not affect transpiration of the mistletoe leaves. Hence, the severity of drought stress for the host is greatly increased by the presence of mistletoes. It may therefore be concluded that the hypothesis tested in the current experiments is valid, i.e. that the flux of ABA from the host xylem into the mistletoe is strongly controlled by the mistletoe and ensures a favourable water and nutritional status for the mistletoe, even if the water status of the host is impaired. However, the present experiments cannot exclude the possibility that this mechanism of control is located at the level of ABA breakdown [30] before it enters the mistletoe leaves, rather than at the level of ABA influx at the haustorium. However such a breakdown in the apoplastic space seems to be unlikely.

Despite the lack of short-term sensitivity of ABA levels in mistletoe leaves to even strong changes of ABA levels in the xylem sap of the host, ABA levels in mistletoe leaves were




Fig. 6. ABA concentrations in leaves ( $\mu\text{mol g dw}^{-1}$ ) of mistletoe leaves after 20 h perfusion of the host xylem with different ABA concentrations. Significant differences in RSA between treatments (ABA levels) were not observed.

relatively high (Fig. 6) compared to ABA levels in the leaves of several tree species, including poplar [1,11,14,40]. This was also observed for the hemiparasite *Rhinanthus* in relation to its host *Hordeum vulgare* [30,31]. Significant transpiration of the mistletoe leaves was maintained (Fig. 1) indicating a diminished sensitivity of the stomata of mistletoe leaves to ABA. Such a diminished sensitivity may have been caused by external pH values unfavourable for ABA action on stomatal aperture [46,45], and/or low calcium and high potassium concentrations, previously reported for mistletoes in general and *Viscum* in particular [6,17,22,33,34,39,44]. High calcium levels play an important role in the signalling cascade [2,25,49] leading to stomatal closure and, thus, for ABA sensitivity of this process [25,48]; high potassium levels may counteract the response on stomatal aperture mediated by calcium [15,48]. Thus, in addition to an effective control of short-term changes in ABA levels, relative ABA insensitivity seems to mediate high transpiration by mistletoe leaves that is largely independent of environmental changes.

The adaxial stomatal density of *Viscum* leaves of  $89 \pm 23$  stomata per mm determined in the present study is lower than those reported in a study performed at the end of the 19th century (100 per  $\text{mm}^2$  adaxial and abaxial) [36]. One possible explanation is that stomatal densities of leaves of *V. album* also decrease with  $\text{CO}_2$  enrichment in the atmosphere as previously reported for other plants [8,55] in order to reduce their expense of transpirational water loss [42,55]. For xylem-tapping mistletoes, for which it is vital to have a more negative water potential than hosts, the reduction of the stomatal density seems to be absurd. But it could make sense if (i) a transpirational gradient is established mainly by the insensitivity of mistletoe stomata at higher VPD [17], or if (ii) temperate mistletoe establishes the difference in water potential mainly osmotically [5,51] or if (iii), due to the increased WUE the host is less water stressed [5] and the relative water potential gradient to the host could be maintained with lower stomatal densities and transpiration rates [9]. Investigations of historical mistletoe material would provide additional evidence on stomatal densities.

We did not analyse stomatal densities of the host leaves. But since previously for *P. x euamericana*, a stomatal density of 170–180 stomata per mm has been reported [50] and other *Populus* species exhibited stomatal densities from 170 to 288 stomata per mm [41], it is likely that the stomatal density of the *V. album* leaves in our study is also considerably lower than those of its host. This would be in agreement with the findings on a range of New Zealand mistletoes, which have in general lower stomatal densities in leaves than its hosts [4]. Further studies have to elucidate whether the mistletoe/host quotients of the stomatal densities in general and particularly for *Viscum* are related to their heterotrophic carbon gain and the quotients mistletoe/host of the photosynthetic capacity.

#### Acknowledgments

The authors thank R.K. Monson for valuable comments on an earlier draft of the manuscript and the Deutsche

Forschungsgemeinschaft (Re 515/12) as well as the Helmholtz-Gemeinschaft within the virtual institute “Stable Isotope Analysis in Ecosystem Research”.

## References

- [1] K. Aasamaa, A. Sober, W. Hartung, U. Ninemets, Rate of stomatal opening, shoot hydraulic conductance and photosynthetic characteristics in relation to leaf abscisic acid concentration in six temperate deciduous trees, *Tree Physiol.* 22 (2002) 267–276.
- [2] C.J. Atkinson, T.A. Mansfield, A.M. Kean, W.J. Davies, Control of stomatal aperture by calcium in isolated epidermal tissue and whole leaves of *Commelina communis* L. *New Phytol.* 111 (1989) 9–17.
- [3] J.T. Ball, I.E. Woodrow, J.A. Berry, A model predicting stomatal conductance and its contribution to the control of photosynthesis under different environmental conditions, in: J. Biggins, M. Nijhoff (Eds.), Dordrecht, 1987, pp. 221–224.
- [4] P. Bannister, G.L. Strong, Stomatal densities and frost resistances of some New Zealand mistletoes, *N.Z. Bot. Soc. Newslett.* 50 (1997) 1–13.
- [5] P. Bannister, G.L. Strong, Carbon and nitrogen isotope ratios, nitrogen content and heterotrophy in New Zealand mistletoes, *Oecologia* 126 (2001) 10–20.
- [6] M. Barberaki, S. Kintzios, Accumulation of selected macronutrients in mistletoe tissue cultures: effect of medium composition and explant source, *Sci. Hortic.* 95 (2002) 133–150.
- [7] W.L. Bauerle, J.E. Toler, G.G. Wang, Stomatal conductance of *Acer rubrum* ecotypes under varying soil and atmospheric water conditions: predicting stomatal responses with an abscisic acid-based model, *Tree Physiol.* 24 (2004) 805–811.
- [8] D.J. Beerling, C.K. Kelly, Stomatal density responses of temperate woodland plants over the past seven decades of CO<sub>2</sub> increase: a comparison of Salisbury (1927) with contemporary data, *Am. J. Bot.* 84 (1997) 1572–1583.
- [9] D.J. Beerling, F.I. Woodward, Ecophysiological responses of plants to global environmental-change since the last glacial maximum, *New Phytol.* 125 (1993) 641–648.
- [10] S.L. Chen, J.K. Li, S.S. Wang, A. Huttermann, A. Altman, Salt, nutrient uptake and transport, and ABA of *Populus euphratica*: a hybrid in response to increasing soil NaCl, *Trees Struct. Funct.* 15 (2001) 186–194.
- [11] S.L. Chen, S.S. Wang, A. Altman, A. Huttermann, Genotypic variation in drought tolerance of poplar in relation to abscisic acid, *Tree Physiol.* 17 (1997) 797–803.
- [12] S.L. Chen, S.S. Wang, A. Huttermann, A. Altman, Xylem abscisic acid accelerates leaf abscission by modulating polyamine and ethylene synthesis in water-stressed intact poplar, *Trees-Struct. Funct.* 16 (2002) 16–22.
- [13] P.J. Clark, F.C. Evans, Distance to nearest neighbor as a measure of spatial relationship in populations, *Ecology* 35 (1954) 445–453.
- [14] H. Cochard, M. Ridolfi, E. Dreyer, Water stress in an ABA-unresponsive hybrid poplar (*Populus koreana* × *trichocarpa* cv. Peace) – response, *New Phytol.* 134 (1996) 455–461.
- [15] M.J. Correia, M.L. Rodrigues, M.L. Osorio, M.M. Chaves, Effects of growth temperature on the response of lupin stomata to drought and abscisic acid, *Aust. J. Plant Physiol.* 26 (1999) 549–559.
- [16] W.J. Davies, J.H. Zhang, Root signals and the regulation of growth and development of plants in drying soil, *Annu. Rev. Plant. Physiol. Plant. Mol. Biol.* 42 (1991) 55–76.
- [17] M.A. El Sharkawy, J.H. Cock, A.D. Hernandez, Differential response of stomata to air humidity in the parasitic mistletoe (*Phthirusa pyrifolia*) and its host, mandarin orange (*Citrus resitilata*), *Photosynth. Res.* 9 (1986) 333–343.
- [18] P. Escher, M. Eiblmeier, I. Hetzger, H. Rennenberg, Seasonal and spatial variation of reduced sulphur compounds in mistletoes (*Viscum album*) and the xylem sap of its hosts (*Populus X euramericana* and *Abies alba*), *Physiol. Plant.* 117 (2003) 72–78.
- [19] P. Escher, M. Eiblmeier, I. Hetzger, H. Rennenberg, Seasonal and spatial variation of carbohydrates in mistletoes (*Viscum album*) and the xylem sap of its hosts (*Populus x euamericana* and *Abies alba*), *Physiol. Plant.* 120 (2004) 212–219.
- [20] P. Escher, M. Eiblmeier, I. Hetzger, H. Rennenberg, Spatial and seasonal variation in amino compounds in the xylem sap of a mistletoe (*Viscum album*) and its hosts (*Populus* spp. and *Abies alba*), *Tree Physiol.* 24 (2004) 639–650.
- [21] P. Escher, M. Eiblmeier, H. Rennenberg, Differences in the influx of glutamine and nitrate into *Viscum album* from the xylem sap of its hosts, *Plant Physiol. Biochem.* 42 (2004) 739–744.
- [22] J.T. Fisher, Water relations of mistletoes and their hosts, in: M. Calder, P. Bernhardt (Eds.), *The Biology of Mistletoes*, Academic Press, Sydney, 1983, pp. 161–184.
- [23] O. Härtel, Über den Wasserhaushalt von *Viscum album* L. *Ber. Dtsch. Bot. Ges.* 55 (1937) 310–321.
- [24] P. Hertz, Über den gegenseitigen durchschnittlichen Abstand von Punkten, die mit bekannter mittlerer Dichte im Raume angeordnet sind, *Math. Ann.* 67 (1909) 387–398.
- [25] A.M. Hetherington, R.S. Quatrano, Mechanisms of action of abscisic acid at the cellular-level, *New Phytol.* 119 (1991) 9–32.
- [26] J. Hintze, NCSS 2004 and PASS, Number Cruncher Statistical Systems, Kaysville, Utah, 2005, Available from: <http://www.ncss.com>.
- [27] J.F. Hodgson, Aspects of Carbon Nutrition of Angiospermous Parasites. Ph.D. thesis, Department of Botany, University of Sheffield, UK, 1973.
- [28] K.R. Hultine, J.D. Marshall, A comparison of three methods for determining the stomatal density of pine needles, *J. Exp. Bot.* 52 (2001) 369–373.
- [29] B. Ihl, F. Jacob, A. Meyer, G. Sembdner, Investigations on the endogenous levels of abscisic acid in a range of parasitic phanerogams, *J. Plant Growth Regul.* 5 (1987) 191–205.
- [30] F. Jiang, W.D. Jeschke, W. Hartung, Water flows in the parasitic association *Rhinanthus minor*/*Hordeum vulgare*, *J. Exp. Bot.* 54 (2003) 1985–1993.
- [31] F. Jiang, W.D. Jeschke, W. Hartung, Abscisic acid (ABA) flows from *Hordeum vulgare* to the hemiparasite *Rhinanthus minor* and the influence of infection on host and parasite abscisic acid relations, *J. Exp. Bot.* 55 (2004) 2323–2329.
- [32] H. Kaiser, Die stomatare Reaktion von *Sambucus nigra* und *Aegopodium podagraria* in Abhängigkeit von Licht und Luftfeuchte, Ph.D. thesis, Mathematisch-Naturwissenschaftliche Fakultät der Christian-Albrechts-Universität zu Kiel, 1999.
- [33] S. Kintzios, M. Barberaki, J. Drossopoulos, P. Turgelis, J. Konstas, Effect of medium composition and explant source on the distribution profiles selected micronutrients in mistletoe tissue cultures, *J. Plant Nutr.* 26 (2003) 369–397.
- [34] B. Lamont, Mineral nutrition of mistletoes, in: M. Calder, P. Bernhardt (Eds.), *The Biology of Mistletoes*, Academic Press, Sydney, 1983, pp. 185–204.
- [35] K. Langer, V. Levchenko, J. Fromm, D. Geiger, R. Steinmeyer, S. Lautner, P. Ache, R. Hedrich, The poplar K<sup>+</sup> channel KPT1 is associated with K<sup>+</sup> uptake during stomatal opening and bud development, *Plant J.* 37 (2004) 828–838.
- [36] G. Marktanner-Turneretscher, Zur Kenntnis des anatomischen Baues unserer Loranthaceen, *Sitzungsber. kaiserl. Akad. Wissensch. XCI* (1885) 1–12.
- [37] R.J. Mertens, B. Deus-Neumann, E.W. Weiler, Monoclonal antibodies for the detection and quantification of the endogenous plant growth regulator, abscisic acid, *FEBS Lett.* 160 (1985) 269–272.
- [38] B. Mühr, 2004. Available from: <http://www.klimadiagramme.de/Bawue/karlsruhe.html>.
- [39] T. Nicoloff, Contribution a la physiologie de la nutrition des parasites végétaux supérieurs, *Revue Gen. Bot.* 35 (1923) 545–552.
- [40] U. Niinemets, A. Sober, O. Kull, W. Hartung, J.D. Tenhunen, Apparent controls on leaf conductance by soil water availability and via light-acclimation of foliage structural and physiological properties in a mixed deciduous, temperate forest, *Int. J. Plant Sci.* 160 (1999) 707–721.
- [41] D.W. Pearce, S. Millard, D.F. Bray, S.B. Rood, Stomatal characteristics of riparian poplar species in a semi-arid environment, *Tree Physiol.* 26 (2006) 211–218.
- [42] J. Penuelas, R. Matamala, Changes in N and S leaf content, stomatal density and specific leaf-area of 14 plant-species during the last 3 centuries of CO<sub>2</sub> increase, *J. Exp. Bot.* 41 (1990) 1119–1124.

- [43] A.D. Peuke, W.D. Jeschke, W. Hartung, The uptake and flow of C, N and ions between roots and shoots in *Ricinus communis* L. III. Long-distance transport of abscisic acid depending on nitrogen nutrition and salt stress, *J. Exp. Bot.* 45 (1994) 741–747.
- [44] M. Popp, A. Richter, Ecophysiology of xylem-tapping mistletoes, *Prog. Bot.* 59 (1998) 659–672.
- [45] L. Prokic, Z. Jovanovic, M.R. Mcainsh, Z. Vucinic, R. Stikic, Species-dependent changes in stomatal sensitivity to abscisic acid mediated by external pH, *J. Exp. Bot.* 57 (2006) 675–683.
- [46] L. Prokic, Z. Jovanovic, R. Stikic, Z. Vucinic, The mutual effect of extracellular  $\text{Ca}^{2+}$ , abscisic acid, and pH on the rate of stomatal closure, In: *Biophysics from Molecules to Brain: in Memory of Radoslav K. Andjus*, vol. 1048 (2005) pp. 513–516.
- [47] W. Rasband, ImageJ, 1.34s, U.S. National Institutes of Health, Bethesda, Maryland, U.S.A., 2006, Available from: <http://rsb.info.nih.gov/ij/>.
- [48] M. Ridolfi, M.L. Fauveau, P. Label, J.P. Garrec, E. Dreyer, Water stress in an ABA-unresponsive hybrid poplar (*Populus koveana* x *trichocarpa* cv. Peace) – response, *New Phytol.* 134 (1996) 445–454.
- [49] M.R.G. Roelfsema, R. Hedrich, In the light of stomatal opening: new insights into ‘the Watergate’, *New Phytol.* 167 (2005) 665–691.
- [50] R.T.W. Siegwolf, R. Matyssek, M. Saurer, S. Maurer, M.S. Gunthardt-Goerg, P. Schmutz, J.B. Bucher, Stable isotope analysis reveals differential effects of soil nitrogen and nitrogen dioxide on the water use efficiency in hybrid poplar leaves, *New Phytol.* 149 (2001) 233–246.
- [51] G.L. Strong, P. Bannister, Water relations of temperate mistletoes on various hosts, *Funct. Plant Biol.* 29 (2002) 89–96.
- [52] F. Tardieu, W.J. Davies, Stomatal response to abscisic-acid is a function of current plant water status, *Plant Physiol.* 98 (1992) 540–545.
- [53] C.v. Tubeuf, *Monographie der Mistel*, Druck und Verlag von R. Oldenbourg, 1923.
- [54] I. Ullmann, O.L. Lange, H. Ziegler, J. Ehleringer, E.D. Schulze, J.R. Cowan, Diurnal courses of leaf conductance and transpiration of mistletoes and their hosts in Central Australia, *Oecologia* 67 (1985) 577–587.
- [55] F.I. Woodward, C.K. Kelly, The influence of  $\text{CO}_2$  concentration on stomatal density, *New Phytol.* 131 (1995) 311–327.